

external link (opens in a new window)

Search Sources Analytics Alerts My list Settings Live Chat Help Tutorials

## Quick Search

Search

Back to results | &lt; Previous 23 of 186 Next &gt;

[Link to Full Text](#) | [Download](#) [Export](#) [Print](#) [E-mail](#) [Create bibliography](#) [Add to My List](#)

## Acta Horticulturae

Volume 739, 2007, Pages 405-410

## Evaluation of ISO method in saffron qualification (Conference Paper)

Hadizadeh, F.<sup>a</sup>, Mahdavi, M.<sup>a</sup>, Emami, S.A.<sup>a</sup>, Khashayarmanesh, Z.<sup>a</sup>, Hassanzadeh, M.<sup>a</sup>, Asili, J.<sup>a</sup>, Seifi, M.<sup>b</sup>, Nassirli, H.<sup>b</sup>, Shariatmoghadam, A.<sup>c</sup>, Noorbakhsh, R.<sup>d</sup>

<sup>a</sup> School of Pharmacy, Mashhad University of Medical Sci., Mashhad, Iran<sup>b</sup> Biotechnology and Pharmaceutical, Sciences Research Center, Mashhad University of Medical Sci., Mashhad, Iran<sup>c</sup> Novin Saffron, P.O. Box 91735-1356, Mashhad, Iran<sup>d</sup> Standard Institute, Mashhad, Iran

## Abstract

[View references \(13\)](#)

The factors which are very important in saffron are crocin (color), picrocrocin (flavor) and safranal (aroma). Higher amount of these compounds in saffron provide higher quality of saffron. ISO (the international organization for standardization) has set a classification of saffron based on minimum requirements of each quality. According to ISO picrocrocin, safranal and crocin are expressed as direct reading of the absorbance of 1% aqueous solution of dried saffron at 257, 330 and 440 nm respectively. There was some doubt about the accuracy of the safranal result using ISO method. Therefore, it was decided to evaluate the amounts of safranal and crocin in saffron according to ISO method and compare the results with the amounts of these compounds when using high performance liquid chromatography (HPLC). In HPLC method in an isocratic run a 30 cm ODS column were used. Mobile phase was mixture of acetonitrile and water (76% v/v). A UV detector at wavelengths of 308 nm for safranal and 440 nm for crocin was also employed. Results indicated that standard crocin had significant absorption at 308 nm. We concluded that present ISO method is not an accurate method of measuring the amount of safranal compounds in saffron. Therefore, it is suggested to evaluate this method and meanwhile using a HPLC method to evaluate the amount of safranal and crocin in saffron samples.

## Reaxys Database Information

|

## Author keywords

Crocin; HPLC; ISO; Picrocrocin; Safranal

ISSN: 05677572 Source Type: Book series Original language: English

Document Type: Conference Paper

## References (13)

[View in table layout](#)
[Page](#) [Export](#) [Print](#) [E-mail](#) [Create bibliography](#)

- Alonso, G.L., Salinas, M.R., Garijo, J.  
1 Methods to determination authenticity of aroma of saffron (*Crocus sativus* L.) (1998) *J. Food Prot.* 61, pp. 1521-1530.  
[Link to Full Text](#)
- Alonso, G.L., Salinas, M.R., Sánchez-Fernández, M.A., Garijo, J.  
2 **Note. Safranal content in Spanish saffron** (2001) *Food Science and Technology International*, 7 (3), pp. 225-229. Cited 12 times.  
[Link to Full Text](#)
- Caballero-Ortega, H., Pereda-Miranda, R., Riverón-Negrete, L., Hernández, J.M., Medécigo-Ríos, M., Castillo-Villanueva, A., Abdullaev, F.I.  
3 Chemical composition of saffron (*Crocus sativus* L.) from four countries (2004) *Acta Hort.* 650, pp. 321-326. Cited 5 times.

## Cited by since 1996

This article has been cited **10 times** in Scopus: (Showing the 2 most recent)

Boskabady, M.H., Tabatabaee, A., Byrami, G.  
**The effect of the extract of *Crocus sativus* and its constituent safranal, on lung pathology and lung inflammation of ovalbumin sensitized guinea-pigs** (2012) *Phytomedicine*

Boskabady, M.H., Seyedhosseini Tamijani, S.M., Rafatpanah, H.  
**The effect of crocus sativus extract on human lymphocytes' cytokines and T helper 2/T helper 1 balance** (2011) *Journal of Medicinal Food*

[View details of all 10 citations](#)

Inform me when this document is cited in Scopus:

[Set alert](#) | [Set feed](#)

## Related documents

Showing the 2 most relevant related documents by all shared references:

Caballero-Ortega, H., Pereda-Miranda, R., Abdullaev, F.I.  
**HPLC quantification of major active components from 11 different saffron (*Crocus sativus* L.) sources** (2007) *Food Chemistry*

Lage, M., Cantrell, C.L.  
**Quantification of saffron (*Crocus sativus* L.) metabolites crocins, picrocrocin and safranal for quality determination of the spice grown under different environmental Moroccan conditions** (2009) *Scientia Horticulturae*

[View all related documents based on all shared references](#) or [select the shared references to use](#)

Find more related documents in Scopus based on:

[Authors](#) | [Keywords](#)

## More By These Authors

The authors of this article have a total of **92 records** in Scopus: (Showing 5 most recent)

Moalem, S.A., Hadizadeh, F., Farimani, E.S., Rameshrad, M., Allahyari, M.  
**Synthesis and immunomodulation of human lymphocyte proliferation and cytokine (interferon- $\gamma$ ) production of four novel leflunomide analogues** (2012) *Medicinal Chemistry*

Emami, S.A., Asgary, S., Naderi, G.A., Ardekani, M.R.S., Aslani, S., Airin, A., Kasher, T., Sahebkar, A.

[Add apps](#) | [Help](#)