

external link (opens in a new window)

Search Sources Analytics Alerts My list Settings Live Chat Help Tutorials

## Quick Search

Search

Back to results | &lt; Previous 150 of 186 Next &gt;

[Link to Full Text](#) | [View at publisher](#) | [Download](#) | [Export](#) | [Print](#) | [E-mail](#) | [Create bibliography](#) | [Add to My List](#)

Iranian Journal of Public Health

Volume 36, Issue 1, 2007, Pages 55-61

## Tc1 cells percentage in patients with cutaneous Leishmaniasis before and after treatment with glucantime

Mohajery, M.<sup>a</sup>, Shamsian, A.A.<sup>a</sup>, Mahmoodi, M.<sup>b</sup><sup>a</sup> Dept. of Medical Parasitology and Mycology, Ghaem Hospital, Mashhad University of Medical Sciences, Mashhad, Iran<sup>b</sup> Bu.Ali Research Inst., Mashhad University of Medical Sciences, Mashhad, Iran

## Abstract

[View references \(27\)](#)

Background: Various responses and different prognosis to specific treatment in different patients from one hand, and importance of IFN- $\gamma$  producer cells on the other hand impressed us to study Tc1. Methods: The study was conducted in Ghaem Medical Center and Bu-Ali Research Institute, Mashhad University of Medical Sciences, Iran from 2001 to 2002. Lymphocytes of 36 patients were counted and cultured. Percentage of different responsible immunity cells in 29 patients, were determined by Flow Cytometry System before and after medication with glucantime (IM). Patients who showed improvement after the treatment were put into group 1 and those who did not recover were labeled group 2. In this self-control clinical trial, sampling method was consecutive non-probability and the results were analyzed by t-test consequently. Results: The percentage of Tc1 cells showed a significant increase despite of being stimulated with Phorpol-Mristate-Acetate (PMA) among the whole studied patients and group 1 ( $P=0.069$  and  $P=0.040$ , respectively). While no significant change was observed among patients in group 2. Conclusion: This verifies the influence of Tc1 cells for the treatment of patients with CL and perhaps the role of glucantime in improving the cell immunity response through increasing such cells.

## Reaxys Database Information

|

## Author keywords

Cutaneous leishmaniasis; Flow cytometry; Glucantime and lymphocyte; Tc1

ISSN: 03044556 Source Type: Journal Original language: English

Document Type: Article

## References (27)

[View in table layout](#)[Add apps](#) | [Help](#)
[Page](#) | [Export](#) | [Print](#) | [E-mail](#) | [Create bibliography](#)

- (2002)
- World Health Organization (updated). Division of Control of Tropical Diseases. Leishmaniasis control. available from [www.who.int/health-topics/leishmaniasis](http://www.who.int/health-topics/leishmaniasis)  
[Link to Full Text](#)
  - Hadighi, R., Mohebbi, M., Boucher, P., Hajjarian, H., Khamesipour, A., Ouellette, M. **Unresponsiveness to glucantime treatment in Iranian cutaneous Leishmaniasis due to drug-resistant Leishmania tropica parasites** (2006) PLoS Medicine, 3 (5), pp. 659-667. Cited 61 times.  
[http://medicine.plosjournals.org/archive/1549-1676/3/5/pdf/10.1371\\_journal.pmed.0030162-L.pdf](http://medicine.plosjournals.org/archive/1549-1676/3/5/pdf/10.1371_journal.pmed.0030162-L.pdf)  
doi: 10.1371/journal.pmed.0030162  
[Link to Full Text](#) | [View at publisher](#)
  - Ajdary, S., Alimohammadian, M.H., Eslami, M.B., Kemp, K., Kharazmi, A.

## Cited by since 1996

This article has been cited 0 times in Scopus.

Inform me when this document is cited in Scopus:

[Set alert](#) | [Set feed](#)

## Related documents

Showing the 2 most relevant related documents by all shared references:

Rogers, K.A., Titus, R.G.  
**The human cytokine response to Leishmania major early after exposure to the parasite in vitro** (2004) *Journal of Parasitology*

Lehmann, J., Alber, G.  
**Murine leishmaniasis: A paradigm for the importance of T helper 1 and T helper 2 cells** (1998) *OIE Revue Scientifique et Technique*

[View all related documents based on all shared references](#) or [select the shared references to use](#)

Find more related documents in Scopus based on:

[Authors](#) | [Keywords](#)

## More By These Authors

The authors of this article have a total of 3 records in Scopus: (Showing 3 most recent)

Mehdikhani Karimabad, H., Shabestari, M., Baharvand, H., Vosough, A., Gourabi, H., Shahverdi, A., Shamsian, A., Abdolhoseini, S., Moazzami, K., Moradi Marjanimehr, M., Emami, F., Bidkhori, H.R., Hamedanchi, A., Talebi, S., Farrokhi, F., Jabbari-Azad, F., Fadavi, M., Garivani, U., Mahmoodi, M., Aghdami, N.  
**Lack of beneficial effects of granulocyte colony-stimulating factor in patients with subacute myocardial infarction undergoing late revascularization: A double-**