

Traumatic hip dislocation in an infant

Peivandi, M.T.^a, Kachooei, A.R.^a, Amel-Farzad, S.^b

^a Department of Orthopedics, **Mashhad University of Medical Sciences, Mashad, Iran**

^b Boo-Ali Center of Research, **Mashhad University of Medical Sciences, Mashad, Iran**

[View references \(1 0\)](#)

Abstract

A ۱۹ month-old child with pain and deformity in hip following a downfall was admitted to the Emergency Department of Emdadi Hospital. After clinical examination and plain X-ray, hip dislocation was diagnosed and the patient was transferred to the emergency operation theater. Under general anesthesia, the hip was simply reduced using closed method. In the following day, a hip spica-cast was applied for three weeks. Regarding the low-energy trauma required in this kind of dislocation and long-term and hazardous complications due to the delayed diagnosis, it is important to consider the probability of hip dislocation in children. © Iranian Red Crescent Medical Journal.

Reaxys Database Information

Author keywords

Close reduction; Dislocation; Hip

Indexed Keywords

EMTREE medical terms: article; bone radiography; case report; cast application; child; clinical examination; emergency ward; falling; femur fracture; fracture reduction; hip dislocation; hip injury; human; preschool child; range of motion

ISSN: ۱۰۶۱۴۳۹۰ **Source Type:** Journal **Original language:** English

Document Type: Article